

Quasi-Phase Transition and Many-Spin Kondo Effects in Graphene Nanodisk

Motohiko Ezawa

Department of Applied Physics, University of Tokyo, Hongo 7-3-1, 113-8656, Japan

The trigonal zigzag nanodisk with size N has N localized spins. We investigate its thermodynamical properties with and without external leads. Leads are made of zigzag graphene nanoribbons or ordinary metallic wires. There exists a quasi-phase transition between the quasi-ferromagnet and quasi-paramagnet states, as signaled by a sharp peak in the specific heat and in the susceptibility. Lead effects are described by the many-spin Kondo Hamiltonian. A new peak emerges in the specific heat. Furthermore, the band width of free electrons in metallic leads becomes narrower. By investigating the spin-spin correlation it is argued that free electrons in the lead form spin-singlets with electrons in the nanodisk. They are indications of many-spin Kondo effects.

Introduction: Graphene nanostructure[1] has the potential for future application in nanoelectronics and spintronics. In particular, much attention is now focused on graphene nanoribbons[2] due to almost flat low-energy band at the Fermi level depending on the edge states. Another basic element of graphene derivatives is a graphene nanodisk[3, 4, 5, 6, 7, 8]. It is a nanometer-scale disk-like material which has a closed edge. There are many type of nanodisks, among which the trigonal zigzag nanodisk is prominent in its electronic and magnetic properties because there exist N -fold degenerated half-filled zero-energy states when its size is N (Fig.1).

In this paper we explore thermodynamical properties of the trigonal zigzag nanodisk. The system is well approximated by the infinite-range Heisenberg model. It is exactly solvable. A sharp peak emerges in the specific heat and in the susceptibility, which we interpret as a quasi-phase transition between the quasi-ferromagnet and quasi-paramagnet states. We then investigate a nanodisk-lead system, where the lead is made of a zigzag graphene nanoribbon or an ordinary metallic wire. We refer to it as a graphene lead or a metallic lead.

It is shown that lead effects are described by the many-spin Kondo Hamiltonian. Electron spins in the nanodisk and the lead orient into the opposite directions to lower the coupling energy. A new peak appears around a certain temperature ($T = T_K$) in the specific heat but not in the susceptibility for small size nanodisks. The internal energy decreases near the peak, and the band width of free electrons in the lead becomes narrower in the instance of the metallic lead. Furthermore, the spin-spin correlation takes the maximum value at $T = 0$, remains almost constant for $T \lesssim T_K$, and then decreases monotonically as T increases. We interpret these phenomena to mean that free electrons in the lead are consumed to make spin-singlets with electrons in the nanodisk. It is intriguing that all electrons (only a few portion of electrons) in the nanodisk are engaged in the case of the graphene (metallic) lead. They are indications of Kondo effects due to the Kondo coupling between electrons in the lead and the nanodisk.




FIG. 1: (a) Trigonal zigzag nanodisks. The size parameter N is defined in this way. The number of carbon atoms is given by $N_C = N^2 + 6N + 6$. (b) The nanodisk-lead system. The nanodisk with $N = 7$ is connected to the right and left leads.

Quasi-Ferromagnet: The size- N zigzag trigonal nanodisk has N -fold degenerated zero-energy states[4], where the gap energy is as large as a few eV. Hence it is a good approximation to investigate the electron-electron interaction physics only in the zero-energy sector, by projecting the system to the subspace made of those zero-energy states. The zero-energy sector consists of N orthonormal states $|f_\alpha\rangle$, $\alpha = 1, 2, \dots, N$, together with the $SU(N)$ symmetry. Let $U_{\alpha\beta}$ and $J_{\alpha\beta}$ be the Coulomb energy and the exchange energy between electrons in the states $|f_\alpha\rangle$ and $|f_\beta\rangle$. It follows[7] that $J_{\alpha\beta} \approx U_{\alpha\beta}$ and that all $J_{\alpha\beta}$ are of the same order of magnitude for any pair of α and β , implying that the $SU(N)$ symmetry is broken but not so strongly. It is a good approximation to start with the exact $SU(N)$ symmetry, by replacing $U_{\alpha\beta}$ and $J_{\alpha\beta}$ with their averages U and J , respectively. Then, the zero-energy sector is described by the Hamiltonian[9],

$$H_D = -JS_{\text{tot}}^2 + \frac{1}{2}U'n_{\text{tot}}^2 + \left(\frac{U}{2} + J\right)n_{\text{tot}}, \quad (1)$$

with $U' \equiv U - \frac{1}{2}J$ and $J \approx U$, where $\mathbf{S}_{\text{tot}} = \sum_\alpha \mathbf{S}(\alpha)$ is the total spin, and $n_{\text{tot}} = \sum_\alpha n(\alpha)$ is the total electron number. Here, $n(\alpha) = d_\sigma^\dagger(\alpha)d_\sigma(\alpha)$, and $\mathbf{S}(\alpha) = \frac{1}{2}d_\sigma^\dagger(\alpha)\boldsymbol{\tau}_{\sigma\sigma'}d_{\sigma'}(\alpha)$ with $d_\sigma(\alpha)$ the annihilation operator of electron with spin $\sigma = \uparrow, \downarrow$ in the state $|f_\alpha\rangle$: $\boldsymbol{\tau}$ is the Pauli matrix. We call $\mathbf{S}_{\text{tot}}$ the nanodisk spin.

Apart from an irrelevant constant, the Hamiltonian (1) is the infinite-range Heisenberg model, $H_S = -JS_{\text{tot}}^2$, in the half-filled sector with $n_{\text{tot}} = N$. The nanodisk spin




FIG. 2: Thermodynamical properties of the nanodisk without leads. (a) The specific heat C in unit of $k_B N$. (b) The entropy S in unit of $k_B N \log 2$. (c) The magnetization $\langle \mathbf{S}_{\text{tot}}^2 \rangle$ in unit of S_g^2 . (d) The susceptibility χ in unit of S_g . The size is $N = 1, 2, 2^2, \dots, 2^{10}$. The horizontal axis stands for the temperature T in unit of JN/k_B . The arrow represents the phase transition point T_c in the limit $N \rightarrow \infty$.

takes the maximum value $S_g = \sqrt{N/2(N/2+1)}$ in the ground state, where all spins are spontaneously polarized. The nanodisk spin system exhibits a strong ferromagnetic order due to a large exchange interaction. The relaxation time is finite but quite large even if the size N is small. We have called such a system quasi-ferromagnet[4].

Thermodynamical Properties: The infinite-range Heisenberg model is exactly diagonalizable, $H_S|\Psi_S\rangle = E_S|\Psi_S\rangle$, with $E_S = -JS(S+1)$, where S takes half-integer or integer values from $N/2$ down to $1/2$ or 0 , depending on whether N is odd or even. The total degeneracy of the energy level E_S is $(2S+1)g_N(S)$ with $g_N(N/2 - q) = N C_q - N C_{q-1}$.

We have a complete set of the eigenenergies together with their degeneracies. The partition function of the nanodisk with size N is exactly calculable,

$$Z_S = \sum_S (2S+1) g_N(S) e^{-\beta JS(S+1)}. \quad (2)$$

According to the standard procedure we can evaluate the specific heat $C_N(T)$, the entropy $S_N(T)$, the magnetization $\langle \mathbf{S}_{\text{tot}}^2 \rangle$ and the susceptibility $\chi = \frac{1}{k_B T} (\langle \mathbf{S}_{\text{tot}}^2 \rangle - \langle \mathbf{S}_{\text{tot}} \rangle^2)$ from this partition function. The entropy is given by $S_N(0) = k_B \log(N+1)$ at zero temperature, corresponding to the ground state multiplicity $N+1$. We display them in Fig.2 for size $N = 1, 2, 2^2, \dots, 2^{10}$.

There appear singularities in thermodynamical quantities as $N \rightarrow \infty$, which represent a phase transition at $T_c \equiv JN/2k_B$ between the ferromagnet and paramagnet states (Fig.2). For finite N , there are steep changes around T_c , though they are not singularities. It is not a phase transition. However, it would be reasonable to call it a quasi-phase transition between the quasi-ferromagnet

and quasi-paramagnet states. Such a quasi-phase transition is manifest even in finite systems with $N = 100 \sim 1000$.

The specific heat and the magnetization take nonzero-values for $T > T_c$ [Fig.2(a),(c)], which is zero in the limit $N \rightarrow \infty$. The entropy for $T > T_c$ is lower than that of the paramagnet [Fig.2(b)]. These results mean the existence of some correlations in the quasi-paramagnet state. The maximum value of the susceptibility increases linearly as N becomes large. It is an indicator of the quasi-phase transition.

Many-Spin Kondo Hamiltonian: We proceed to investigate how thermodynamical properties of the nanodisk is affected by the attachment of the leads. Though there are two leads attached to a nanodisk, the lead Hamiltonian H_L and the transfer Hamiltonian H_T are expressed as if there were a single lead after a certain transformation[7],

$$H_L = \sum_{k\sigma} \varepsilon(k) c_{k\sigma}^\dagger c_{k\sigma}, \quad (3a)$$

$$H_T = \tilde{t} \sum_{k\sigma} \sum_{\alpha} \left(c_{k\sigma}^\dagger d_{\sigma}(\alpha) + d_{\sigma}^\dagger(\alpha) c_{k\sigma} \right), \quad (3b)$$

where $c_{k\sigma}$ is the annihilation operator of electron in the lead with the wave number k and the dispersion relation $\varepsilon(k)$.

When charges transfer between the nanodisk and the leads, the total electron number n_{tot} is no longer fixed in the nanodisk. However, the nanodisk remains to be half-filled, when a charge transfers from the lead to the nanodisk and then transfers back from the nanodisk to the lead. The process is the second order effect in the tunneling coupling constant $\tilde{t}$. We derive the effective Hamiltonian for such a process.

The total Hamiltonian is $H = H_D + H_L + H_T$. We take $H_0 = H_D + H_L$ as the unperturbed term and H_T as the perturbation term. Note that $U \gg \tilde{t}$. We make a canonical transformation known as the Schrieffer-Wolff transformation[10] to eliminate H_T , $H \rightarrow \tilde{H} = e^{iG} H e^{-iG}$, with G the generator satisfying $H_T + \frac{i}{2}[G, H_0] = 0$. The dominant contribution comes from the Fermi surface, $\varepsilon(k) = \varepsilon_F$. We assume the symmetric condition $\varepsilon_F = U + \frac{1}{2}U'$ with respect to the Fermi energy. Then, after a straightforward calculation, we obtain $\tilde{H} = H_D + H_L + H_K + O(\tilde{t}^3)$, where H_K is the second order term in $\tilde{t}$. It is the many-spin Kondo Hamiltonian,

$$H_K = J_K \sum_{kk'\sigma\sigma'} c_{k\sigma}^\dagger \tau_{\sigma\sigma'} c_{k'\sigma'} \cdot \mathbf{S}_{\text{tot}}, \quad (4)$$

with the Kondo coupling constant $J_K = 8\tilde{t}^2/U'$. The difference between the above many-spin Kondo Hamiltonian and the ordinary Kondo Hamiltonian is whether the local spin is given by the summation over many spins $\mathbf{S}_{\text{tot}}$ or a single spin $\mathbf{S}$. Note that $\mathbf{S}_{\text{tot}}^2$ is a dynamical variable but $\mathbf{S}^2$ is not, $\mathbf{S}^2 = 3/4$.




FIG. 3: Thermodynamical properties of the nanodisk (a,b) with graphene leads and (a',b') with metallic leads. See the caption of Fig.2. We have set $J_K/J = 0.2$ and $D = 2k_B T_c$. A new peak appears around T_K in the specific heat.

Functional Integration: The total Hamiltonian is now given by $H_{\text{eff}} = H_S + H_L + H_K$ at half filling. We define the spinor $\psi = (c_\uparrow, c_\downarrow)^t$. The partition function in the Matsubara form is given in terms of the Hamiltonian density $\mathcal{H}_{\text{eff}}$ as

$$Z = \text{Tr}_S \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \exp \left[- \int_0^\beta d\tau \int dx (\psi^\dagger \partial_\tau \psi + \mathcal{H}_{\text{eff}}) \right] \\ = \text{Tr}_S [\exp(-\beta H_S) Z_K], \quad (5)$$

with $Z_K = \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \exp[-S_K]$, where S_K is the action $S_K = \int_0^\beta d\tau \int dx (\psi^\dagger \partial_\tau \psi + \mathcal{H}_L + \mathcal{H}_K)$. We first perform a functional integral over the lead electron's degree of freedom in Z_K , and then summed up the nanodisk spin in (5).

Because an electron in the lead is constrained within a very narrow region, it is a good approximation to neglect momentum scatterings,

$$H_K \simeq 2J_K \mathbf{s} \cdot \mathbf{S}_{\text{tot}}, \quad (6)$$

where $\mathbf{s} = \frac{1}{2} \sum_{k\sigma\sigma'} c_{k\sigma}^\dagger \tau_{\sigma\sigma'} c_{k\sigma'}$ is the electron spin in the lead. The action S_K is summarized as

$$S_K = \int \frac{d\omega}{2\pi} \sum_k \psi^\dagger(k) M(k) \psi(k), \quad (7)$$

with $M(k) = -[i\omega - \varepsilon(k)] + J_K \tau \cdot \mathbf{S}_{\text{tot}}$. Performing the functional integration we find $Z_K = \text{Det}[M] = \exp[-\beta F_K]$, where F_K is the Helmholtz free energy,

$$F_K = -\frac{1}{2\beta} \sum_k \ln [\cosh(\beta J_K |\mathbf{S}_{\text{tot}}|) + \cosh(\beta \varepsilon(k))]. \quad (8)$$

This is reduced to the well-known formula for free electrons with the dispersion relation $\varepsilon(k)$ for $J_K = 0$.

Zigzag Graphene Nanoribbon Leads: We first consider the system where the leads are made of zigzag graphene nanoribbons. Owing to the flat band at the zero energy, $\varepsilon(k) = 0$, the result of the functional integration (8) is quite simple,

$$F_K = -\frac{1}{\beta} \ln \cosh[\beta J_K |S_{\text{tot}}|/2]. \quad (9)$$

The effective Hamiltonian for the nanodisk spin is $H_S + F_K$. The lead effect is to make the effective spin stiffness larger and the ferromagnet more rigid.

The trace over the nanodisk spin is carried out in (5),

$$Z = \sum_S (2S+1) g_N(S) e^{-\beta J S(S+1)} \cosh \left(\frac{\beta J_K}{2} \sqrt{S(S+1)} \right). \quad (10)$$

We compare thermodynamical properties of the nanodisk with leads (Fig.3) and without leads (Fig.2). The magnetization $\langle \mathbf{S}_{\text{tot}}^2 \rangle$ and the susceptibility χ are found to be indistinguishable from those of the nanodisk without leads (Fig.2). In Fig.3, we show the specific heat $C_G(T)$ and the entropy $S_G(T)$ for various size N . The significant feature is the appearance of a new peak in the specific heat at $T = T_K \approx (J_K/2J)T_c$, though it disappears for large N . We examine the internal energy $E_G(T)$, which is found to decrease around T_K (Fig.4). Near zero temperature it reads

$$E_G(T) \simeq -J S_g^2 - \frac{J_K}{2} S_g + J_K S_g e^{-\beta J_K S_g}. \quad (11)$$

The first term ($\propto J$) represents the energy stabilization due to the ferromagnetic order present in the nanodisk without leads, while the second term ($\propto J_K$) represents the one due to the Kondo coupling. Furthermore, it follows that the entropy is reduced at zero temperature as $S_G(0) - S_N(0) = -k_B \log 2$, as implies that the ground state multiplicity at the zero temperature is just one half of that of the system without leads.

The spin-spin correlation $|\langle \mathbf{s} \cdot \mathbf{S}_{\text{tot}} \rangle|$ is calculated based on the partition function (5) and shown in Fig.5(a). Near zero temperature we find

$$|\langle \mathbf{s} \cdot \mathbf{S}_{\text{tot}} \rangle| \simeq \frac{1}{2} S_g \tanh[\beta J_K S_g/2]. \quad (12)$$

It takes the maximum value $S_g/2$ at $T = 0$, and remains almost constant for $T \lesssim T_K$, and then monotonically decreases as T increases. Finally, it almost vanishes in the quasi-paramagnet phase for large N since $\langle \mathbf{S}_{\text{tot}} \rangle \approx 0$. We may interpret these phenomena as follows. Electrons in the lead and the nanodisk form spin-singlet states to lower the coupling energy (6). The singlet state is rather tight for $T \lesssim T_K$, but thermally broken as T increases.

All these features indicate the occurrence of the Kondo effect due to the coupling between the spins in the nanodisk and the leads.




FIG. 4: The internal energy E in unit of Nk_B for the nanodisk (a) without leads and (b) with graphene leads. See the caption of Fig.2. (a) The energy decreases except for $N = 1$ as the temperature decreases, which represents the ferromagnetic order. (b) There exists an additional energy decrease around T_K , which is prominent for $N = 1$, attributed to the Kondo effect.




FIG. 5: The spin-spin correlation in unit of $S_g/2$. See the caption of Fig.3. The correlation occurs due to the Kondo coupling. It is almost constant for $T \lesssim T_K$, and decreases as T increases.

Metallic Leads: Next we consider the system comprised of metallic leads with a constant energy density, $\rho(\varepsilon) = \rho$ for $|\varepsilon| < D$ and $\rho(\varepsilon) = 0$ for $|\varepsilon| > D$. We change the momentum integration into the energy integration in (8),

$$F_K = -\frac{\rho}{2\beta} \int_{-D}^D d\varepsilon \ln [\cosh \beta J_K |\mathbf{S}_{\text{tot}}| + \cosh \beta \varepsilon]. \quad (13)$$

The free energy is given by $F_K = -\rho\beta^{-2} [\beta^2 D J_K |\mathbf{S}_{\text{tot}}| - 2\beta D \log 2 + \text{Li}_2(-e^{\beta(D-J_K|\mathbf{S}_{\text{tot}}|)}) - \text{Li}_2(-e^{-\beta(D+J_K|\mathbf{S}_{\text{tot}}|)})]$, where $\text{Li}_2[x]$ is the dilogarithm function[11]. It is reduced to the free energy of the nanodisk with graphene leads in the limit $D \rightarrow 0$ with $\rho = 1/2D$.

The magnetization $\langle \mathbf{S}_{\text{tot}}^2 \rangle$ and the susceptibility χ are found to be indistinguishable from those in the case of no leads (Fig.2). The relation $S_M(0) - S_N(0) = -k_B \log 2$ holds for the entropy precisely as in the case of graphene leads. In Fig.3, we show the specific heat $C_M(T)$ and the entropy $S_M(T)$ for various size N . A broad peak appears at $T = T_K$ in the specific heat. Using the asymptotic behaviors[11] of $\text{Li}_2[x]$, we obtain the free energy in low

temperature regime as

$$F_K \simeq \frac{-\rho}{\beta} \left[\beta D J_K |\mathbf{S}_{\text{tot}}| - 2D \log 2 + \frac{\pi^2}{6\beta} + \frac{\beta}{2} (D - J_K |\mathbf{S}_{\text{tot}}|)^2 \right], \quad (14)$$

which is valid up to the terms in the order of $e^{-\beta D}$. The entropy $S_M(T)$, the specific heat $C_M(T)$ and the internal energy $E_M(T)$ read as follows,

$$S_M(T) \simeq k_B \log \frac{N+1}{2} + \frac{\pi^2}{3} \rho k_B^2 T, \quad C_M(T) \simeq \frac{\pi^2}{3} \rho k_B^2 T, \quad (15)$$

and $E_M(T) = E_G(T) + \Delta E(T)$ with

$$\Delta E(T) \simeq -\frac{\rho}{2} (D - J_K S_g)^2 + \frac{\pi^2}{6} \rho (k_B T)^2. \quad (16)$$

All terms proportional to ρ have arisen from free electrons in the metallic lead. The internal energy $E_M(T)$ consists of two terms: $E_G(T)$ is identical to the energy (11) for the nanodisk with graphene leads, and $\Delta E(T)$ is the energy of the metallic lead. The first term of $\Delta E(T)$ shows that the band width of free electrons in the lead becomes narrower due to the Kondo coupling. We may interpret that n free electrons with

$$n = \rho J_K S_g \quad (17)$$

are consumed to make spin-coupling with electrons in the nanodisk. The second term is the thermal energy of free electrons in the metallic lead.

We show the spin-spin correlation $|\langle \mathbf{s} \cdot \mathbf{S}_{\text{tot}} \rangle|$ in Fig.5(b). The overall features are the same as for the nanodisk with graphene leads. However, there are new features. First of all, the value of correlation is quite small. This is because spin-singlets are formed only by a small portion of electrons in the metallic lead which are near the Fermi level. We expect that this number density is given by (17). Indeed, it is observed that $|\langle \mathbf{s} \cdot \mathbf{S}_{\text{tot}} \rangle| \rightarrow \frac{1}{2} \rho J_K S_g$ at $T = 0$ as $N \rightarrow \infty$ [Fig.5(b)].

In this paper we have investigated thermodynamical properties of a zigzag graphene nanodisk without and with leads. The lead effects are summarized by the many-spin Kondo Hamiltonian. One effect is to enhance the ferromagnetic order. This result is important to manufacture spintronic circuits by connecting leads in nanodevices[9]. We have shown various thermodynamical results indicating many-spin Kondo effects.

I am very much grateful to N. Nagaosa for many fruitful discussions on the subject. This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture No. 20840011.

-
- [1] K. S. Novoselov, *et al.*, Science **306**, 666 (2004). K. S. Novoselov, *et al.*, Nature **438**, 197 (2005). Y. Zhang, *et al.*, Nature **438**, 201 (2005).

- [2] M. Fujita, *et al.*, J. Phys. Soc. Jpn. **65**, 1920 (1996). M. Ezawa, Phys. Rev. B, **73**, 045432 (2006). L. Brey, and H. A. Fertig, Phys. Rev. B, **73**, 235411 (2006). F. Muñoz-Rojas, *et al.*, Phys. Rev. B, **74**, 195417 (2006). Y. -W Son, M. L. Cohen, and S. G. Louie, Phys. Rev. Lett., **97**, 216803 (2006). V. Barone, O. Hod, and G. E. Scuseria, Nano Lett., **6**, 2748 (2006). M. Y. Han, *et al.*, Phys. Rev. Lett., **98**, 206805 (2007).
- [3] M. Ezawa, Physica Status Solidi (c) **4**, No.2, 489 (2007).
- [4] M. Ezawa, Phys. Rev. B **76**, 245415 (2007); M. Ezawa, Physica E **40**, 1421-1423 (2008).
- [5] J. Fernández-Rossier, and J. J. Palacios, Phys. Rev. Lett. **99**, 177204 (2007).
- [6] O. Hod, V. Barone, and G. E. Scuseria, Phys. Rev. B **77**, 035411 (2008).
- [7] M. Ezawa, Phys. Rev. B **77**, 155411 (2008).
- [8] W. L. Wang, S. Meng and E. Kaxiras, Nano Letters **8**, 241 (2008). W. L. Wang, *et al.*, Phys. Rev. Lett. **102**, 157201 (2009).
- [9] M. Ezawa, Eur. Phys. J. B **67**, 543 (2009)
- [10] J. R. Schrieffer and P. A. Wolff, Phys. Rev. **149**, 491 (1966).
- [11] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, New York: Dover, 1004-1005, (1972).